



A Review of the Relationship Between Vitamin D and Parkinson Disease Symptoms

Michelle E. Fullard^{1*} and John E. Duda^{2,3}

¹ Department of Neurology, University of Colorado, Denver, CO, United States, ² Parkinson's Disease Research, Education and Clinical Center, Corporal Michael J. Crescenzo VA Medical Center, Philadelphia, PA, United States, ³ Department of Neurology, University of Pennsylvania School of Medicine, Philadelphia, PA, United States

OPEN ACCESS

Edited by:

Amie Hiller,

Oregon Health and Science University,
United States

Reviewed by:

Patrick McCullough,

Wright State University, United States

Chien Tai Hong,

Taipei Medical University, Taiwan

*Correspondence:

Michelle E. Fullard

michelle.fullard@cuanschutz.edu

Specialty section:

This article was submitted to
Movement Disorders,
a section of the journal
Frontiers in Neurology

Received: 27 January 2020

Accepted: 28 April 2020

Published: 27 May 2020

Citation:

Fullard ME and Duda JE (2020) A
Review of the Relationship Between
Vitamin D and Parkinson Disease
Symptoms. *Front. Neurol.* 11:454.
doi: 10.3389/fneur.2020.00454

Vitamin D is a fat-soluble secosteroid that exerts its effects by binding to the vitamin D receptor (VDR), through which it directly and indirectly modulates the expression of hundreds to thousands of genes. While originally known for its role in regulating calcium homeostasis and metabolism, vitamin D is now associated with many other health conditions, including Parkinson's disease (PD). A high prevalence of vitamin D deficiency has been noted in PD for at least the past two decades. These findings, along with the discovery that the VDR and 1 α -hydroxylase, the enzyme that converts vitamin D to its active form, are highly expressed in the substantia nigra, led to the hypothesis that inadequate levels of circulating vitamin D may lead to dysfunction or cell death within the substantia nigra. Studies investigating the relationship between vitamin D status and PD, however, have been inconsistent. Two prospective studies examined the association between mid-life vitamin D levels and risk of PD and produced conflicting results—one demonstrated an increased risk for PD with lower mid-life vitamin D levels, and the other showed no association between vitamin D and PD risk. One of the most consistent findings in the literature is the inverse association between serum vitamin D level and motor symptom severity in cross-sectional studies. While these data suggest that vitamin D may modify the disease, another likely explanation is confounding due to limited mobility. Fall risk has been associated with vitamin D in PD, but more study is needed to determine if supplementation decreases falls, which has been demonstrated in the general population. The association between vitamin D and non-motor symptoms is less clear. There is some evidence that vitamin D is associated with verbal fluency and verbal memory in PD. Studies in PD have also shown associations between vitamin D status and mood, orthostatic hypotension and olfactory impairment in PD. While more research is needed, given the numerous potential benefits and limited risks, vitamin D level assessment in PD patients and supplementation for those with deficiency and insufficiency seems justified.

Keywords: vitamin D, Parkinson's disease, motor symptoms, review, non-motor symptoms

INTRODUCTION

Vitamin D refers to a group of fat-soluble secosteroids that can be obtained through dermal synthesis by exposure to sunlight or through dietary intake (1). Vitamin D itself is biologically inert and must undergo two hydroxylations to become active. The liver converts vitamin D to 25-hydroxyvitamin D and then the kidney and other tissues convert 25-hydroxyvitamin D to the active form 1,25-dihydroxyvitamin D, also known as calcitriol (1). Vitamin D exerts its effects by binding to the vitamin D receptor (VDR), through which it directly and indirectly modulates the expression of hundreds to thousands of genes (1, 2). While originally known for its role in regulating calcium homeostasis and metabolism, vitamin D was reported in the 1930s and 1940s to improve conditions like psoriasis, asthma and rheumatoid arthritis (3, 4). It is now associated with many other health conditions, such as cardiovascular disease, cancer, autoimmune disease and neurodegenerative disease, including Parkinson's disease (PD) (5).

Multiple mechanisms have been proposed for the role of vitamin D in neurodegenerative diseases. Vitamin D may confer neuroprotection through the action of neurotrophic factors, regulation of nerve growth or through protection against cytotoxicity (6). In several studies, the synthesis of neurotrophic factors, including neurotrophin 3 (NT3) and glial cell line-derived neurotrophic factor (GDNF), were upregulated by 1,25(OH)₂D₃ (7–12). Riaz et al. and Wang et al. showed that vitamin D induced stimulation of neurotrophins was correlated with a neuroprotective effect in rat models (13, 14). Through the VDR, Vitamin D can upregulate nerve growth factor (NGF) (7, 15), and in a rat model of peripheral nerve injury, rats exposed to vitamin D₂ demonstrated significantly increased axogenesis and axon diameters (16).

Vitamin D may also modulate the toxicity of reactive oxygen species. Nitric oxide is produced by inducible nitric oxide synthase (iNOS), an enzyme that is induced in CNS neurons and non-neuronal cells as part of an immune response (17). At high levels, nitric oxide can damage neurons, and vitamin D has been shown to inhibit the synthesis of iNOS, thereby reducing levels of nitric oxide (17, 18). In hippocampal neurons, the presence of vitamin D was associated with downregulation of L-type voltage-sensitive Ca²⁺ channel and was correlated with a neuroprotective effect at physiologic doses (19). Lastly, vitamin D may exert a neuroprotective effect through reducing oxidative stress. In a study in which dopaminergic and non-dopaminergic neurons were exposed to glutamate, pretreatment with vitamin D protected both cell types against cytotoxicity in a dose-dependent manner (20).

The high prevalence of vitamin D deficiency in Parkinson's disease was first noted by Sato and colleagues in 1997 (21). Since then, multiple studies have demonstrated a higher prevalence of vitamin D deficiency and insufficiency in PD compared to controls (22–24). After the discovery that the VDR and 1 α -hydroxylase, the enzyme that converts vitamin D to its active form, were highly expressed in the substantia nigra, it was hypothesized that inadequate levels of circulating vitamin D may lead to dysfunction or cell death within the substantia

nigra (25, 26). Adding further support to this hypothesis were VDR knockout mice that demonstrated motor impairment and reduced stride length (27), and animal and cell culture studies that demonstrated neuroprotective effects of vitamin D against MPTP and 6-OHDA exposure (28–30). Over the last two decades, multiple studies have examined the relationship between low levels of vitamin D and Parkinson's disease. In this manuscript, we will review the evidence for the effect of hypovitaminosis D on Parkinson's disease risk, symptoms and progression.

PARKINSON'S DISEASE RISK

Over the past decade, several studies have attempted to determine the relationship between hypovitaminosis D and the risk of Parkinson's disease, with conflicting results (Table 1). The first study to examine the association between vitamin D levels in mid-life and the risk of Parkinson's disease was performed using a nationwide drug-reimbursement database in Finland (31). Of the 3,173 men and women included in the study, 50 went on to develop Parkinson's disease over the 29-year follow-up period. After adjusting for potential confounders, participants in the highest quartile for baseline serum vitamin D level had a 65% lower risk of developing PD than those in the lowest quartile, suggesting that lower levels of vitamin D in mid-life may increase the risk of Parkinson's disease (31). In a case control study in elderly Chinese, self-reported outdoor activity and total vitamin D intake were inversely associated with Parkinson's disease, suggesting vitamin D may be protective against the development of PD (32). However, recall bias and the lack of serum vitamin D levels make it difficult to draw firm conclusions from this study. In addition, outdoor physical activity was used as a proxy for vitamin D status, but there is compelling data that physical activity reduces the risk of Parkinson's disease, which is a likely confounder in this study (33).

More recent data have failed to demonstrate an association between vitamin D levels and Parkinson's disease risk. In contrast to the study in Finland, an analysis from the prospective Atherosclerosis Risk in Communities (ARIC) study cohort of almost 13,000 men and women in the United States found no association between baseline vitamin D level and risk of PD 17 years later (34). In a smaller study of participants at high risk for Parkinson's disease in the Parkinson Associated Risk Syndrome (PARS) study, there was no difference in baseline serum vitamin D level between the high risk group (dopamine transporter scan deficit and hyposmia) vs. the lower risk group (35). Taken together, these more recent data do not support the hypothesis that chronic vitamin D insufficiency contributes to PD pathogenesis. All of these studies, however, were based on a single serum measurement of vitamin D. To try to overcome some of these limitations, Larsson et al. performed a mendelian randomization study to determine if genetically decreased vitamin D concentrations are associated with PD (36). They used 4 single-nucleotide polymorphisms (SNPs) associated with serum vitamin D level and found no association between these genetic variants and PD (36). The authors noted that

TABLE 1 | Association of Vitamin D with Parkinson's disease risk and symptoms.

	Study findings	References
Parkinson's Disease Risk		
Vitamin D and Parkinson's Disease Risk	Findings are mixed. A study in Finland found an association between vitamin D level and future PD risk while a study in the United States did not. Differences in the populations studied may play a role	(31–36)
Vitamin D Receptor and Parkinson's Disease Risk	The SNP FokI is associated with a decreased risk of PD in Asian populations but not in Caucasian populations	(37–40)
Motor Symptoms		
Disease Severity	Lower serum vitamin D levels are consistently associated with higher motor symptom severity in PD, however, reduced mobility in advanced disease may result in limited sun exposure and lower vitamin D levels	(22, 41–47)
Disease Progression	Findings are mixed. In two studies, lower baseline vitamin D level was a predictor of increased motor severity, while two other studies found no association. There is risk for confounding due to limited mobility and sun exposure as the disease progresses	(22, 48–53)
Balance and Falls	Lower vitamin D levels are associated with falls. High dose vitamin D supplementation may improve balance in younger PD patients. More recent data in the general population suggests there is a U-shaped response curve to vitamin D supplementation	(45, 54–75)
Non-motor Symptoms		
Cognition	In PD without dementia, higher plasma vitamin D levels are associated with better verbal fluency and verbal memory. The VDR FokI AA genotype is associated with faster cognitive decline and lower VDR activity	(44, 50, 76, 77)
Mood	There is limited evidence of an association between depression and lower serum vitamin D levels in PD. Data is inconsistent in the general population, however, a meta-analysis of 31,000 participants found lower vitamin D levels in participants with depression compared to controls	(44, 73, 78–84)
Orthostatic hypotension	Lower serum vitamin D levels were associated with orthostatic hypotension in one study involving PD patients and multiple studies involving the general population, however, the findings have been inconsistent	(85–90)
Olfactory function	One study in PD reported a negative correlation between odor identification scores and serum vitamin D level. More studies are needed	(91)

UPDRS, Unified Parkinson's Disease Rating Scale; H&Y, Hohn and Yahr; SNP, Single Nucleotide Polymorphism.

these four SNPs only explain about 3.6% of the variation in 25-hydroxyvitamin D concentrations, which would limit the power to detect a weaker association between these SNPs and PD (36).

Potential explanations for the inconsistent findings between the two main prospective studies include differences in geography, behaviors, and economics of the populations studied (92). Vitamin D levels in the Finnish population were much lower than in the US population, so it is possible vitamin D levels below a certain threshold are required to increase the risk of PD. In addition, diet and activity levels may differ between the populations, both of which are associated with vitamin D levels and PD risk (93). Given the differences in geography, environmental risk factors for PD may differ as well.

Multiple meta-analyses have examined the association between VDR polymorphisms and risk of Parkinson's disease (37–40). The most recent and comprehensive meta-analysis by Wang et al. included the SNPs FokI, TaqI, ApaI, and Bsm and included 5–7 studies per SNP with over one thousand PD patients and controls (37). They found that FokI was significantly associated with decreased risk of PD in most models (allelic, dominant, and additive) but not in a recessive model. However, when stratified by race, FokI was associated with PD susceptibility in all four models in Asian populations while there was no significant association in any of the models in the Caucasian population. TaqI was associated with an increased risk of PD in a dominant model only, and there were no significant associations between ApaI and Bsm in any model. In addition,

on sensitivity analysis, omission of any one of five studies for TaqI resulted in a lack of association (37). More work is needed in this area with larger numbers and more diverse populations.

MOTOR SYMPTOMS

Disease Severity

The motor symptoms of Parkinson's disease include bradykinesia, rigidity, rest tremor and postural instability. Symptom severity can be measured using several scales, including the Unified Parkinson's Disease Rating Scale (UPDRS) (41), as well as the Hoehn and Yahr stage (H&Y) (42). One of the most consistent findings in the literature is the association between serum vitamin D levels and motor symptom severity in PD. In multiple cross-sectional studies, vitamin D level was correlated with the motor portion of the UPDRS (22, 43–47), and H&Y stage (43). Suzuki et al. found a significant, inverse relationship between serum vitamin D level and part III of the UPDRS in 137 PD participants (43). Similarly, vitamin D level was highest in participants with H&Y stage 1–1.5 (mean 23.9 ng/ml) and lowest in participants who were H&Y stages 4–5 (mean 14.5 ng/ml) and demonstrated a dose-response relationship (43). A second study using data from 388 PD patients in the Harvard Biomarker Study had similar findings. Total UPDRS score was inversely correlated with serum vitamin D level ($p = 0.02$) after adjusting for age, sex, race and vitamin D supplementation (22). In this study, however, there was no association between vitamin D level and H&Y stage.

While these data, along with preclinical studies in mice and cell culture (28–30), suggest that vitamin D modifies the disease, another very plausible explanation is confounding due to mobility. Increasing disease severity is associated with a decline in mobility, which leads to decreased sun exposure and decreased dermal synthesis of vitamin D. Therefore, the lower vitamin D levels documented in more advanced PD may be a result of limited sun exposure and represent reverse causation, rather than an effect of lower vitamin D levels on PD severity.

Associating genotype and clinical outcome is one way to try to avoid reverse causality. The evidence involving vitamin D receptor (VDR) polymorphisms and Parkinson's disease risk and severity, however, has been inconsistent (37). In a study by Suzuki, the FokI CC genotype for VDR polymorphism was associated with a milder form of PD, but vitamin D level was not (43). In an intervention study performed by the same group, 114 PD patients were randomized to receive either 1,200 units of vitamin D3 daily or placebo for 12 months (94). The H&Y stage was stable in the treatment group, but significantly declined in the placebo group. FokI genotype modified the relationship between vitamin D supplementation and motor progression. For participants with the FokI CC genotype, vitamin D supplementation did not affect progression (compared to the placebo group), but Vitamin D supplementation prevented deterioration in those with FokI TT (94). This genotype has been shown to have less transcriptional activity (76), and therefore individuals with this genotype may require higher serum levels of vitamin D to perform the same functions as those with the CC genotype. More study in larger populations is needed to determine if this is a stable finding.

Progression

Four studies have evaluated the relationship between disease progression and serum vitamin D level in Parkinson's disease and one study has evaluated the relationship between vitamin D supplementation and disease progression (Table 1). Evatt et al. used data from 157 participants in the DATATOP study to evaluate the prevalence of vitamin D insufficiency in early, untreated PD, as well as examine the association between vitamin D levels and disease progression (48). There were high rates of vitamin D deficiency and insufficiency in the PD participants, but vitamin D level unexpectedly increased by a small amount over the course of the study (median follow-up of 13 months). There was no relationship between vitamin D level and disease progression (48). Similarly, 624 PD patients from Oxford's Parkinson's Disease Centre Discovery cohort had baseline serum vitamin D measured and were followed for a mean of 3.2 years, and no association was found between baseline vitamin D level and disease progression measured by the MDS-UPDRS III (49). However, reduced levels of vitamin D were associated with worse activities of daily living at baseline as measured by MDS-UPDRS II. In a third study, 145 newly diagnosed Parkinson's patients and 94 age-matched controls in the North East of England had serum vitamin D levels checked at baseline and 18 months (50). The vitamin D levels at these two time points were significantly lower in the PD participants compared to controls. Lower baseline vitamin D level was a predictor of increased

motor severity at 36 months as measured by the MDS-UPDRS III (50). Lastly, Ding and colleagues found that lower vitamin D levels at baseline predicted a greater change in UPDRS score at follow-up (12–25 months) after adjusting for age, sex, race and vitamin D supplementation (22). Similar to the cross-sectional studies demonstrating an association between serum vitamin D level and motor severity, these studies on motor progression are also at risk for confounding due to more limited mobility as the disease progresses.

Using data from the National Institutes of Health Exploratory Trials in Parkinson's Disease (NET-PD) Long-term study, Luthra et al. examined disease progression at 3 years in patients with early PD who took 400 IU or more per day of vitamin D compared to those who did not take supplements (51). Of the 1,741 participants, only 12% were taking a vitamin D supplement and 34% were taking a multivitamin that contained vitamin D. At 3 year follow-up, there was no difference in UPDRS scores between those taking vitamin D and those who were not.

Data regarding the effects of vitamin D supplementation on Parkinson's disease progression is sparse. A case report from 1997 described a 50 year old patient with 13 years of PD symptoms somewhat responsive to levodopa therapy, bone abnormalities, low serum calcium and phosphorus and low serum vitamin D (52). Vitamin D3 supplementation at 4,000 IU daily of 25-hydroxyvitamin D3, roughly equivalent to 20,000 IU of vitamin D3 (53), and calcium supplements were added to his regimen and his lab values normalized. In addition, his parkinsonism significantly improved over the following year and he was able to reduce his levodopa by half (52). While this is an extreme case, it highlights the need for more study on the effects of vitamin D supplementation on PD symptoms. To date, only two small randomized, placebo-controlled trials have been done examining the effects of vitamin D supplementation on motor symptoms and progression in PD. As discussed above, Suzuki et al. randomized 114 PD patients to receive 1,200 IU/day of vitamin D3 or placebo for 12 months. The serum vitamin D level doubled for those in the intervention group and remained unchanged for the placebo group. H&Y stage was stable in the intervention group, while H&Y stage significantly worsened for the placebo group (+0.33) (94). In a second study in Iran, 120 PD patients with levodopa-induced dyskinesia were randomized to receive either 1,000 IU of vitamin D per day or placebo (95). At 3 month follow-up, there was no difference in levodopa-induced dyskinesia as measured by the UPDRS IV sub score and no difference in the UPDRS motor score, suggesting that vitamin D has no effect on dyskinesia. The follow-up time, however, was too short to determine if vitamin D supplementation influenced motor progression and the dose of vitamin D supplementation was low.

Both of these studies used low doses of vitamin D supplementation, which may have limited the potential to detect improvement in symptoms. A report in 2005 noted that 10–15 min of whole-body exposure to sunlight in the peak of summer can generate 10,000–25,000 IU of vitamin D3, concluding that the recommendations for vitamin D intake of 200–800 IU/day are likely very inadequate (96, 97). A study from 1977 showed that a daily intake of 10,000 IU daily resulted in serum vitamin

D levels most similar to the subjects who were exposed to daily whole-body artificial ultraviolet light (53). One of the reasons often cited for giving lower doses of vitamin D supplementation is the concern for toxicity and hypercalcemia, however, multiple studies have shown that toxicity is rare and high serum levels of vitamin D are not strongly correlated with hypercalcemia (4, 98). When hypercalcemia occurs, it is often easily reversed (4, 98) One state psychiatric hospital in Cincinnati, Ohio offered all 4,700 patients admitted to the facility from 2011 to 2018 supplements of either 5,000 IU/d or 10,000 IU/d of vitamin D3, many of whom were on the supplements for over 12 months (4). Serum 25(OH) D levels were frequently above 100 ng/ml for patients taking doses of 10,000 IU/d, with a range up to 202 ng/ml. None of these patients developed hypercalcemia, nephrolithiasis or other adverse effects and the authors concluded that long-term supplementation at doses of 5,000 to 10,000 U/d appear safe (4). Given the large range of vitamin D supplementation that has been reported in the literature, and the low doses of supplementation provided in the two PD RCTs, more high quality studies with a range of doses are needed to determine if vitamin D is effective for PD motor symptoms and progression and at what doses.

Balance and Falls

Falls are a major determinant of poor quality of life, morbidity and mortality in Parkinson's disease (99, 100). Balance problems and postural instability are the leading cause of falls, with up to 70% of PD patients experiencing at least one fall per year (101–105). Over the past two decades, there has been increasing evidence that vitamin D supplementation can improve falls in older adults, which prompted two studies in the Parkinson's population that evaluated the relationship between vitamin D and balance and falls.

Most of what is known about the association between vitamin D and falls comes from literature in the older, non-PD population. There is a high prevalence of falls among adults ages 65 years and older, which, similar to PD, can lead to fractures and increased morbidity, mortality and cost (106–108). The rate of vitamin D insufficiency is higher in fallers, estimated up to 70%, compared to 40–50% in non-fallers over the age of 65 (54–56). One of the main mechanisms of falls in the elderly is reported to be impairment of postural reactions—either altered or delayed—due to abnormalities in musculoskeletal mechanisms or processing of sensorimotor information (57–59). Low serum vitamin D levels have been associated with increased postural sway (60) and balance (61) and vitamin D supplementation was shown to improve balance and body sway (62) and stabilize postural equilibrium in the elderly (63). In addition, multiple RCTs have shown a decrease in fall rate and fracture rate with vitamin D supplementation in older adults (64–67), especially in those who were vitamin D insufficient at baseline and received at least 800 IU per day (68). In a meta-analysis of five randomized controlled trials of vitamin D supplementation including 1,237 participants, vitamin D supplementation reduced the odds ratio of falling by 22% compared to patients receiving calcium or placebo and further reduced falls in older women by 46–65% (64).

More recent data, however, has suggested that there is a U-shaped response curve when it comes to fall reduction secondary to vitamin D supplementation (69–72). In one study, there was no decrease in falls on low dose vitamin D (400–800 IU), significant decrease in falls on medium doses (1,600–3,200 IU) and no decrease on high doses (4,000 IU) (72). Similarly, when the 12 month serum 25(OH)D levels were divided into quintiles, fall rates also followed a U-shaped curve. The greatest decrease in falls occurred in the group with a final serum 25(OH)D level of 32–38 ng/ml and fall rates increased when the serum 25(OH)D level exceeded 40–45 ng/ml (72). Further trials are needed to determine the appropriate dose of vitamin D supplementation and target serum vitamin D level to reduce falls.

Patients with Parkinson's disease are more likely to fall than their age-matched peers and falls in this population can have devastating consequences (109). While falls have been associated with vitamin D level in Parkinson's disease (73), few studies have examined the mechanisms related to these falls, and only one RCT has examined the effects of vitamin D supplementation on balance. In a study of forty PD patients with a mean H&Y stage of 2.6, serum vitamin D levels were correlated with symmetry of automatic postural responses measured by the Motor Control Test (MCT), which assesses the subject's ability to recover after an unexpected external disturbance (74). In the PD participants, lower vitamin D concentration was correlated with a higher degree of asymmetry in the postural response after adjusting for motor severity (74). In this study, the authors also measured postural sway and found that postural sway was correlated with vitamin D serum levels in only one out of six conditions (45). To determine if vitamin D supplementation improved balance, similar to non-PD older adults, a pilot RCT was performed which involved 16 weeks of high dose vitamin D (10,000 IU/day) or placebo in 51 PD subjects. The authors found that high dose supplementation was well-tolerated and resulted in a doubling of serum concentrations (75). There was no improvement in the primary endpoint of balance as measured by the Sensory Organization Test (SOT). However, a *post hoc* analysis found that younger participants (ages 52–66 years) had significant improvement in the SOT at 16 weeks compared to the older participants (ages 67–86 years), suggesting a potential role for high dose vitamin D supplementation in younger PD patients (75). Given the more recent data consisting of increased falls with high doses of vitamin D in the non-PD population, more studies with longer follow-up are needed to determine an appropriate dose of vitamin D for PD patients at risk for falls.

NON-MOTOR SYMPTOMS

Cognition

While many studies have demonstrated an association between vitamin D level and cognition in the general population, the few clinical trials that have been done fail to show improvement with supplementation (110–119). Even less is known about the effects of vitamin D on cognition in Parkinson's disease, and study findings have been inconsistent (Table 1).

In a study by Peterson et al, 286 participants with PD underwent neuropsychiatric testing and serum vitamin D testing

at baseline (44). Models were adjusted for age, disease duration and Hoehn and Yahr stage. In the non-demented cohort ($n = 225$), higher plasma vitamin D was associated with better scores on tests of verbal fluency and verbal memory. There was no significant association between cognitive testing and vitamin D level in the smaller ($n = 61$) demented subset (44). A second study by Gatto and colleagues assessed the role of VDR polymorphisms in cognitive decline in patients with PD (77). One hundred ninety PD patients in the Parkinson Environment Gene (PEG) study underwent the Mini-Mental State Exam (MMSE) at baseline and up to three follow-up visits and were genotyped for seven VDR polymorphisms. The FokI polymorphism was associated with decline in the MMSE score over the follow-up period after accounting for age, sex, education, and PD disease duration. Participants with AA genotype experienced faster cognitive decline as measured by change in the MMSE. A subset of study participants underwent a battery of neuropsychiatric tests, and those with the AA genotype tended to perform lower on individual neuropsychiatric tests compared to those with the AG or GG genotype (77). The VDR protein that results from the A variant has less transcriptional activity compared to the G variant allele (76), suggesting that lower VDR activity contributes to cognitive decline.

In a prospective observational study performed in England, 145 newly diagnosed PD patients were followed for 3 years and were assessed for disease severity and cognition as part of the Impairment in Cohorts with Longitudinal Evaluation—Parkinson's disease (ICICLE-PD) (50). Serum vitamin D levels were measured at baseline and 18 months. Baseline vitamin D level was not a predictor of future cognitive impairment in this cohort (50).

Several mechanisms have been proposed for the role of vitamin D deficiency in cognitive decline in the general population that may also apply to the PD population. Vitamin D may be involved in the regulation of acetylcholine and clearing of amyloid beta, both of which are implicated in the pathogenesis of cognitive impairment in PD (120, 121). Many studies demonstrate that the VDR and enzymes involved in D3 metabolism are expressed in the central nervous system, particularly in the areas of hippocampus (122, 123), and animal studies have shown that vitamin D deficiency negatively affects hippocampal learning and memory through gene expression and neural development (124). These studies suggest potential mechanisms for the effect of vitamin D on cognitive impairment in PD, but more work is needed to determine if vitamin D status affects cognition in PD and whether supplementation may help prevent or improve cognitive decline.

Mood

The relationship between vitamin D status and depression has been inconsistent in the general population, with some studies demonstrating an increased risk of depression with lower levels of vitamin D (78, 79), while others have shown no association (80, 81). A meta-analysis of over 31,000 participants found that patients with depression had lower vitamin D levels compared to controls (82). Additionally, when comparing the lowest vitamin D category to the highest in cross-sectional studies, those in the lowest category were 30% more likely

to have depression (82). Few studies, however, have examined the effect of vitamin D supplementation on mood. These include a randomized controlled trial in Norway in which 441 overweight or obese participants received either placebo, 20,000 IU or 40,000 IU of vitamin D weekly for a year (83). There was significant improvement in the Beck Depression Inventory in the two intervention arms at 1 year compared to the placebo (83). An open label study of fifty women with type II diabetes and depressive symptoms were given weekly vitamin D supplementation (50,000 IU of ergocalciferol) for 6 months, and there was a significant decrease in depression and anxiety over the study period (84).

Limited evidence exists for the association between depression and vitamin D concentration in Parkinson's disease. Peterson et al. found a correlation between plasma vitamin D levels and depression scores in 225 non-demented PD patients (44). A more recent study by Zhang et al. included 182 subjects with PD and examined the association between vitamin D levels and various motor and non-motor symptoms. They found that vitamin D level was correlated with depression and anxiety scores after adjusting for age, sex and BMI (73). More research is needed in this area to clarify the relationship between vitamin D status and depression in Parkinson's disease and to determine if supplementation is effective at improving mood.

The underlying mechanism by which vitamin D may affect mood is unknown, but several hypotheses exist based on vitamin D's effect on gene expression. The VDR is present in the cingulate cortex and hippocampus, both of which are involved in the pathophysiology of depression (25). In addition, vitamin D response elements have been detected in the promotor regions of serotonin genes, providing further evidence for a potential role of vitamin D in the development of depression (125).

Orthostatic Hypotension

Orthostatic hypotension is defined by the American Academy of Neurology (AAN) as a reduction in systolic blood pressure by at least 20 mm Hg or diastolic blood pressure by at least 10 mm Hg within 3 min of standing (126). It can be symptomatic if patients experience symptoms of lightheadedness, dizziness, blurred vision, fatigue, nausea, etc., or asymptomatic (126, 127), and is estimated to affect 40–60% of PD patients throughout their disease course (128, 129).

The mechanism by which vitamin D may exert an effect on standing blood pressure is unclear. The cells of blood vessel walls all express the VDR and 1-alpha-hydroxylase, which may allow vitamin D involvement in modulating the vascular response when standing (85, 130). In addition, it has been postulated that vitamin D is a renin-angiotensin system regulator (RAS). RAS is implicated in blood pressure regulation and influences the sympathetic nervous system, therefore a vitamin D associated disruption in RAS could lead to dysfunction of the sympathetic system (87, 131).

Vitamin D status has been associated with orthostatic hypotension in one study involving PD patients and multiple studies involving the general population (85–90). In a study by Jang et al. 55 PD participants were divided into two groups—those with orthostatic hypotension (OH) and those without

(130). The group with OH had significantly lower serum 25-hydroxyvitamin D and calcitriol levels. In addition, systolic and diastolic blood pressures, as well as OH symptom severity, were negatively correlated with serum vitamin D and calcitriol levels (130). Disease duration and disease severity were similar between the two groups, making it less likely that the OH group had lower vitamin D due to greater disease severity and more limited mobility. However, the presence of OH itself affects activity levels and is often correlated with disease severity. In a second study involving PD patients, nocturnal blood pressure changes were used as an indicator of autonomic dysfunction, and vitamin D levels were measured (132). Thirty-five PD patients were classified as “dippers” if there was a decline in mean nighttime blood pressure of >10% (normal finding), “non-dippers” if the decline in mean nighttime blood pressure was <10% (pathological) and “reverse dippers” if there was an increase in mean nighttime blood pressure (pathological). There were no significant differences in vitamin D level among the three groups, however, there were only 4 patients in the “dipper” group, so it is difficult to make firm conclusions (132).

In the general population, multiple smaller studies have shown associations between vitamin D level and orthostatic hypotension. McCarroll et al. demonstrated serum 25(OH)D levels were lower in elderly individuals with orthostatic hypotension compared to those without OH (86). Similarly, women over the age of 80 years were more likely to have diastolic orthostatic hypotension if they were vitamin D deficient (88). In the only longitudinal study that has been done examining vitamin D status and OH, it was found that baseline vitamin D deficiency was associated with an increased risk of developing orthostatic hypotension over 1 year follow-up in a group of 51 subjects from the general population (85). A meta-analysis that included 3,646 participants from five studies confirmed the findings that participants with low vitamin D had about twice the odds of having orthostatic hypotension than those within the normal range of vitamin D, and those with orthostatic hypotension were more likely to have lower vitamin D levels (87).

More recent larger studies, however, did not find a link between vitamin D status and OH. An analysis of data from 4,209 participants in The Irish Longitudinal Study on Aging (TILSA) and 2,640 participants in the Progetto Veneto Anziani (Pro.V.A.) study showed no association between vitamin D deficiency or insufficiency and orthostatic hypotension (131, 133). In a subanalysis of a randomized controlled trial assessing vitamin D supplementation for systolic hypertension, high dose supplementation did not improve orthostatic hypotension over a 12 month period (90). It is not clear why these discrepancies exist, but differences in the populations studied may have contributed. At this time, we have conflicting studies from the general population regarding the relationship between orthostatic hypotension and vitamin D, and only one study supporting the relationship in PD, so further studies are needed.

Olfactory Function

Olfactory impairment is a common non-motor symptom in Parkinson's disease, with a prevalence of 50–90% (134–136). It is often one of the first signs of PD (137), which correlates with early involvement of the olfactory bulb with synuclein

pathology (138). In a study of 39 *de novo* PD patients in South Korea, serum 25(OH)D3 level was associated with odor identification score using sniffin' sticks (91). The serum 25(OH)D3 level was also negatively correlated with subjective olfactory dysfunction. The authors postulate that vitamin D plays a role in the pathogenesis of olfactory dysfunction in PD through several potential mechanisms involving calcitriol, the active form of vitamin D (91). Calcitriol is involved in neuronal cell differentiation and neuroprotection through the VDR (6, 139, 140). A study by Glaser et al. demonstrated that the rat olfactory system contains VDR throughout (141), indicating a potential neuroprotective role for calcitriol through VDR signaling. In addition, alterations in dopamine and acetylcholine signaling have been implicated in olfactory dysfunction in Parkinson's disease (142, 143). Calcitriol has been shown to increase choline acetyltransferase and tyrosine hydroxylase activity, which are enzymes responsible for the synthesis of acetylcholine and dopamine, and may in turn exert a neuroprotective effect (144). Lastly, calcium signaling is thought to play a central role in the function of olfactory receptor neurons and can be affected by vitamin D (145). While several possible mechanisms exist for the effect of vitamin D on olfactory dysfunction in Parkinson's disease, the evidence is preliminary, and more study is needed to make any conclusions.

CONCLUSIONS

The relationship between vitamin D status and Parkinson's disease remains unclear, and whether or not supplementation is beneficial beyond bone health in PD has yet to be determined. While preclinical studies in animals and cell culture have shown promising neuroprotective effects of vitamin D, studies in humans have been inconsistent, likely in part due to differing methodologies and populations studied. The most consistent finding in the literature is a higher prevalence of vitamin D deficiency and insufficiency in PD compared to controls, as well as an inverse relationship between vitamin D level and motor severity, but these findings may be due to reduced mobility and decreased sun exposure as PD progresses rather than disease modification by vitamin D. Much less is known about the relationship between vitamin D levels and non-motor symptoms. However, given the numerous potential benefits and limited risks, vitamin D level assessment in PD patients and supplementation for those with deficiency and insufficiency seems justified.

AUTHOR CONTRIBUTIONS

MF contributed to the concept of the review, performed the literature review, and drafted the manuscript. JD contributed to the concept of the review and revised the manuscript for intellectual content.

FUNDING

MF receives funding from the American Brain Foundation, American Academy of Neurology, the Parkinson's Foundation, and the Michael J. Fox Foundation. JD receives funding

from Rehabilitation Research and Development Service of the Department of Veterans Affairs, RX-B1097-I, PI, 2013-2018,

NIH, U01-NS094340, Co-PI, 2015-2018, and the Michael J. Fox Foundation.

REFERENCES

- Cesari M, Incalzi RA, Zamboni V, Pahor M. Vitamin D hormone: a multitude of actions potentially influencing the physical function decline in older persons. *Geriatr Gerontol Int.* (2011) 11:133–42. doi: 10.1111/j.1447-0594.2010.00668.x
- Shirvani A, Kalajian TA, Song A, Allen R, Charoenngam N, Lewanczuk R, et al. Variable genomic and metabolic responses to varying doses of Vitamin D supplementation. *Anticancer Res.* (2020) 40:535–43. doi: 10.21873/anticancer.13982
- Rappaport BZ, Reed CI, Hathaway ML, Struck HC. The treatment of hay fever and asthma with viosterol of high potency. *J Allergy.* (1934) 5:541–53. doi: 10.1016/S0021-8707(34)90130-1
- McCullough PJ, Lehrer DS, Amend J. Daily oral dosing of vitamin D3 using 5000 TO 50,000 international units a day in long-term hospitalized patients: insights from a seven year experience. *J Steroid Biochem Mol Biol.* (2019) 189:228–39. doi: 10.1016/j.jsbmb.2018.12.010
- Bouillon R. Extra-skeletal effects of vitamin D. *Front Horm Res.* (2018) 50:72–88. doi: 10.1159/000486072
- Garcion E, Wion-Barbot N, Montero-Menei CN, Berger F, Wion D. New clues about vitamin D functions in the nervous system. *Trends Endocrinol Metab.* (2002) 13:100–5. doi: 10.1016/S1043-2760(01)00547-1
- Brown J, Bianco JJ, McGrath JJ, Eyles DW. 1,25-Dihydroxyvitamin D 3 induces nerve growth factor, promotes neurite outgrowth and inhibits mitosis in embryonic rat hippocampal neurons. *Neurosci Lett.* (2003) 343:139–43. doi: 10.1016/S0304-3940(03)00303-3
- Wion D, MacGrogan D, Neveu I, Jehan F, Houlgatte R, Brachet P. 1,25-Dihydroxyvitamin D3 is a potent inducer of nerve growth factor synthesis. *J Neurosci Res.* (1991) 28:110–4. doi: 10.1002/jnr.490280111
- Neveu I, Naveilhan P, Baudet C, Brachet P, Metsis M. 1,25-Dihydroxyvitamin D3 regulates NT-3, NT-4 but not BDNF mRNA in astrocytes. *Neuroreport.* (1994) 6:124–6. doi: 10.1097/00001756-199412300-00032
- Musiol IM, Feldman D. 1,25-dihydroxyvitamin D3 induction of nerve growth factor in L929 mouse fibroblasts: effect of vitamin D receptor regulation and potency of vitamin D3 analogs. *Endocrinology.* (1997) 138:12–8. doi: 10.1210/endo.138.1.4858
- Veenstra TD, Fahnestock M, Kumar R. An AP-1 site in the nerve growth factor promoter is essential for 1,25-dihydroxyvitamin D3-mediated nerve growth factor expression in osteoblasts. *Biochemistry.* (1998) 37:5988–94. doi: 10.1021/bi972965+
- Naveilhan P, Neveu I, Wion D, Brachet P. 1,25-Dihydroxyvitamin D3, an inducer of glial cell line-derived neurotrophic factor. *Neuroreport.* (1996) 7:2171–5. doi: 10.1097/00001756-199609020-00023
- Riaz S, Malcangio M, Miller M, Tomlinson DR. A vitamin D3 derivative (CB1093) induces nerve growth factor and prevents neurotrophic deficits in streptozotocin-diabetic rats. *Diabetologia.* (1999) 42:1308–13. doi: 10.1007/s001250051443
- Wang Y, Chiang YH, Su TP, Hayashi T, Morales M, Hoffer BJ, et al. Vitamin D3 attenuates cortical infarction induced by middle cerebral arterial ligation in rats. *Neuropharmacology.* (2000) 39:873–80. doi: 10.1016/S0028-3908(99)00255-5
- Marini F, Bartoccini E, Cascianelli G, Voccoli V, Baviglia MG, Magni MV, et al. Effect of 1 α ,25-dihydroxyvitamin D3 in embryonic hippocampal cells. *Hippocampus.* (2010) 20:696–705. doi: 10.1002/hipo.20670
- Chabas JF, Alluin O, Rao G, Garcia S, Lavaut MN, Risse JJ, et al. Vitamin D2 potentiates axon regeneration. *J Neurotrauma.* (2008) 1247–56. doi: 10.1089/neu.2008.0593
- Garcion E, Sindji L, Montero-Menei C, Andre C, Brachet P, Darcy F. Expression of inducible nitric oxide synthase during rat brain inflammation: regulation by 1,25-dihydroxyvitamin D3. *Glia.* (1998) 22:282–94.
- Garcion E, Nataf S, Berod A, Darcy F, Brachet P. 1,25-Dihydroxyvitamin D3 inhibits the expression of inducible nitric oxide synthase in rat central nervous system during experimental allergic encephalomyelitis. *Mol Brain Res.* (1997). 45:255–67. doi: 10.1016/S0169-328X(96)00260-4
- Brewer LD, Thibault V, Chen KC, Langub MC, Landfield PW, Porter NM. Vitamin D hormone confers neuroprotection in parallel with downregulation of L-type calcium channel expression in hippocampal neurons. *J Neurosci.* (2001) 21:98–108. doi: 10.1523/JNEUROSCI.21-01-00098.2001
- Ibi M, Sawada H, Nakanishi M, Kume T, Katsuki H, Kaneko S, et al. Protective effects of 1 α ,25-(OH) $_2$ D $_3$ against the neurotoxicity of glutamate and reactive oxygen species in mesencephalic culture. *Neuropharmacology.* (2001) 40:761–71. doi: 10.1016/S0028-3908(01)00009-0
- Sato Y, Kikuyama M, Oizumi K. High prevalence of vitamin D deficiency and reduced bone mass in Parkinson's disease. *Neurology.* (1997) 49:1273–8. doi: 10.1212/WNL.49.5.1273
- Ding H, Dhima K, Lockhart KC, Locascio JJ, Hoising AN, Duong K, et al. Unrecognized vitamin D3 deficiency is common in Parkinson disease: Harvard biomarker study. *Neurology.* (2013) 81:1531–7. doi: 10.1212/WNL.0b013e3182a95818
- Evatt ML, DeLong MR, Khazai N, Rosen A, Triche S, Tangpricha V. Prevalence of vitamin D insufficiency in patients with Parkinson disease and Alzheimer disease. *Arch Neurol.* (2008) 65:1348–52. doi: 10.1001/archneur.65.10.1348
- Zhao Y, Sun Y, Ji HF, Shen L. Vitamin D levels in Alzheimer's and Parkinson's diseases: a meta-analysis. *Nutrition.* (2013) 29:828–32. doi: 10.1016/j.nut.2012.11.018
- Eyles DW, Smith S, Kinobe R, McGrath JJ. Distribution of the vitamin D receptor and 1 α -hydroxylase in human brain. *J Chem Neuroanat.* (2005) 29:21–30. doi: 10.1016/j.jchemneu.2004.08.006
- Newmark HL, Newmark J. Vitamin D and Parkinson's disease - a hypothesis. *Mov Disord.* (2007) 22:461–8. doi: 10.1002/mds.21317
- Burne THJ, McGrath JJ, Eyles DW, Mackay-Sim A. Behavioural characterization of vitamin D receptor knockout mice. *Behav Brain Res.* (2005) 157:299–308. doi: 10.1016/j.bbr.2004.07.008
- Shinpo K, Kikuchi S, Sasaki H, Moriwaka F, Tashiro K. Effect of 1,25-dihydroxyvitamin D3 on cultured mesencephalic dopaminergic neurons to the combined toxicity caused by L-buthionine sulfoximine and 1-methyl-4-phenylpyridine. *J Neurosci Res.* (2000) 62:374–82. doi: 10.1002/1097-4547(20001101)62:3<374::AID-JN7>3.0.CO;2-7
- Smith MP, Fletcher-Turner A, Yurek DM, Cass WA. Calcitriol protection against dopamine loss induced by intracerebroventricular administration of 6-hydroxydopamine. *Neurochem Res.* (2006) 31:533–9. doi: 10.1007/s11064-006-9048-4
- Wang JY, Wu JN, Cherg TL, Hoffer BJ, Chen HH, Borlongan C V., et al. Vitamin D3 attenuates 6-hydroxydopamine-induced neurotoxicity in rats. *Brain Res.* (2001) 904:67–75. doi: 10.1016/S0006-8993(01)02450-7
- Knekt P, Kilkkinen A, Rissanen H, Marniemi J, Sääksjärvi K, Heliövaara M. Serum vitamin D and the risk of Parkinson disease. *Arch Neurol.* (2010) 67:808–11. doi: 10.1001/archneurol.2010.120
- Zhu D, Liu GY, Lv Z, Wen SR, Bi S, Wang WZ. Inverse associations of outdoor activity and vitamin D intake with the risk of Parkinson's disease. *J Zhejiang Univ Sci B.* (2014) 15:923–7. doi: 10.1631/jzus.B1400005
- Fang X, Han D, Cheng Q, Zhang P, Zhao C, Min J, et al. Association of levels of physical activity with risk of Parkinson disease: a systematic review and meta-analysis. *JAMA Netw Open.* (2018) 1:e182421. doi: 10.1001/jamanetworkopen.2018.2421
- Shrestha S, Lutsey PL, Alonso A, Huang X, Mosley TH, Chen H. Serum 25-hydroxyvitamin D concentrations in Mid-adulthood and Parkinson's disease risk. *Mov Disord.* (2016) 31:972–8. doi: 10.1002/mds.26573
- Fullard ME, Xie SX, Marek K, Stern M, Jennings D, Siderowf A, et al. Vitamin D in the Parkinson Associated Risk Syndrome (PARS) study. *Mov Disord.* (2017) 32:1636–40. doi: 10.1002/mds.27127

36. Larsson SC, Singleton AB, Nalls MA, Richards JB. No clear support for a role for vitamin D in Parkinson's disease: a mendelian randomization study. *Mov Disord.* (2017) 32:1249–52. doi: 10.1002/mds.27069
37. Wang X, Shen N, Lu Y, Tan K. Vitamin D receptor polymorphisms and the susceptibility of Parkinson's disease. *Neurosci Lett.* (2019) 370:100–6. doi: 10.1016/j.neulet.2019.02.018
38. Lee YH, Kim JH, Song GG. Vitamin D receptor polymorphisms and susceptibility to Parkinson's disease and Alzheimer's disease: a meta-analysis. *Neurosci Lett.* (2014) 5:1947–53. doi: 10.1007/s10072-014-1868-4
39. Li C, Qi H, Wei S, Wang L, Fan X, Duan S, et al. Vitamin D receptor gene polymorphisms and the risk of Parkinson's disease. *Neurosci Lett.* (2015) 36:247–55. doi: 10.1007/s10072-014-1928-9
40. Niu MY, Wang L, Xie AM, ApaI, BsmI, FokI, and TaqI polymorphisms in the vitamin D receptor gene and Parkinson's disease. *Chin Med J.* (2015) 128:1809–14. doi: 10.4103/0366-6999.159358
41. Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, et al. Movement disorder society-sponsored revision of the unified Parkinson's disease rating scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov Disord.* (2008) 23:2129–70. doi: 10.1002/mds.22340
42. Goetz CG, Poewe W, Rascol O, Sampaio C, Stebbins GT, Counsell C, et al. Movement disorder society task force report on the hoehn and yahr staging scale: status and recommendations. *Mov Disord.* (2004) 19:1020–8. doi: 10.1002/mds.20213
43. Suzuki M, Yoshioka M, Hashimoto M, Murakami M, Kawasaki K, Noya M, et al. 25-hydroxyvitamin D, vitamin D receptor gene polymorphisms, and severity of Parkinson's disease. *Mov Disord.* (2012) 27:264–71. doi: 10.1002/mds.24016
44. Peterson AL, Murchison C, Zabetian C, Leverenz JB, Watson GS, Montine T, et al. Memory, mood, and vitamin d in persons with Parkinson's disease. *J Parkinsons Dis.* (2013) 3:547–55. doi: 10.3233/JPD-130206
45. Peterson-Hiller AL, Lobb BM, Murchison C, Quinn JF. The effects of vitamin D supplementation on balance, motor, and neuropsychiatric function in Parkinson's Disease (PD). *Mov Disord.* (2015) 30:S111. doi: 10.1002/mds.26296
46. Marek K, Jennings D, Lasch S, Siderowf A, Tanner C, Simuni T, et al. The Parkinson Progression Marker Initiative (PPMI). *Prog Neurobiol.* (2011) 95:629–35. doi: 10.1016/j.pneurobio.2011.09.005
47. Chitsaz A, Maracy M, Basiri K, Izadi Boroujeni M, Tanhaei AP, Rahimi M, et al. 25-Hydroxyvitamin D and severity of Parkinson's disease. *Int J Endocrinol.* (2013) 2013:689149. doi: 10.1155/2013/689149
48. Evatt ML, DeLong MR, Kumari M, Auinger P, McDermott MP, Tangpricha V. High prevalence of hypovitaminosis D status in patients with early Parkinson disease. *Arch Neurol.* (2011) 68:314–9. doi: 10.1001/archneurol.2011.30
49. Lawton M, Baig F, Toulson G, Morovat A, Evetts SG, Ben-Shlomo Y, et al. Blood biomarkers with Parkinson's disease clusters and prognosis: the Oxford Discovery Cohort. *Mov Disord.* (2020) 5:279–87. doi: 10.1002/mds.27888
50. Sleeman I, Aspray T, Lawson R, Coleman S, Duncan G, Khoo TK, et al. The role of vitamin d in disease progression in early Parkinson's disease. *J Parkinsons Dis.* (2017) 7:669–75. doi: 10.3233/JPD-171122
51. Luthra NS, Kim S, Zhang Y, Christine CW. Characterization of vitamin D supplementation and clinical outcomes in a large cohort of early Parkinson's disease. *J Clin Mov Disord.* (2018) 5:7. doi: 10.1186/s40734-018-0074-6
52. Derex L, Trouillas P. Reversible parkinsonism, hypophosphoremia, and hypocalcemia under vitamin D therapy: case report. *Mov Disord.* (1997) 12:424. doi: 10.1002/mds.870120424
53. Stamp TCB, Haddad JG, Twigg CA. Comparison of oral 25-hydroxycholecalciferol, vitamin D, and ultraviolet light as determinants of circulating 25-hydroxyvitamin D. *Lancet.* (1977) 1:1341–3. doi: 10.1016/S0140-6736(77)92553-3
54. Stein MS, Wark JD, Scherer SC, Walton SL, Chick P, Di Carantonio M, et al. Falls relate to vitamin D and parathyroid hormone in an Australian nursing home and hostel. *J Am Geriatr Soc.* (1999) 47:1195–201. doi: 10.1111/j.1532-5415.1999.tb05199.x
55. Gloth FM, Tobin JD. Vitamin D Deficiency in older people. *J Am Geriatr Soc.* (1995) 43:822–8. doi: 10.1111/j.1532-5415.1995.tb07059.x
56. Thomas MK, Lloyd-Jones DM, Thadhani RI, Shaw AC, Deraska DJ, Kitch BT, et al. Hypovitaminosis D in medical inpatients. *N Engl J Med.* (1998) 338:777–83. doi: 10.1056/NEJM199803193381201
57. Pai YC, Wening JD, Runtz EF, Iqbal K, Pavol MJ. Role of feedforward control of movement stability in reducing slip-related balance loss and falls among older adults. *J Neurophysiol.* (2003) 90:755–62. doi: 10.1152/jn.01118.2002
58. Covinsky KE, Kahana E, Kercher K, Schumacher JG, Kahana B, Justice AC. History and mobility exam index to identify community-dwelling elderly persons at risk of falling. *J Gerontol Ser A Biol Sci Med Sci.* (2001) 56:M253–9. doi: 10.1093/gerona/56.4.M253
59. Cummings SR, Nevitt MC. A hypothesis: the causes of hip fractures. *J Gerontol.* (1989) 44:M107–11. doi: 10.1093/geronj/44.5.M107
60. Dhesi JK, Bearne LM, Moniz C, Hurley M V., Jackson SHD, Swift CG, et al. Neuromuscular and psychomotor function in elderly subjects who fall and the relationship with vitamin D status. *J Bone Miner Res.* (2002) 17:891–7. doi: 10.1359/jbmr.2002.17.5.891
61. Akdeniz S, Hegguler S, Öztürk C, Atamaz FC. The relation between vitamin D and postural balance according to clinical tests and tetraz posturography. *J Phys Ther Sci.* (2016) 28:1272–7. doi: 10.1589/jpts.28.1272
62. Pfeifer M, Begerow B, Minne HW, Suppan K, Fahrleitner-Pammer A, Dobnig H. Effects of a long-term vitamin D and calcium supplementation on falls and parameters of muscle function in community-dwelling older individuals. *Osteoporos Int.* (2009) 20:315–22. doi: 10.1007/s00198-008-0662-7
63. Dhesi JK, Jackson SHD, Bearne LM, Moniz C, Hurley MV, Swift CG, et al. Vitamin D supplementation improves neuromuscular function in older people who fall. *Age Ageing.* (2004) 33:589–95. doi: 10.1093/ageing/afh209
64. Bischoff-Ferrari HA, Dawson-Hughes B, Willett WC, Staehelin HB, Bazemore MG, Zee RY, et al. Effect of vitamin D on falls: a meta-analysis. *J Am Med Assoc.* (2004) 291:1999–2006. doi: 10.1001/jama.291.16.1999
65. Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck AE, Theiler R, et al. Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ.* (2009) 339:843. doi: 10.1136/bmj.b3692
66. Bischoff HA, Stäehelin HB, Dick W, Akos R, Knecht M, Salis C, et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res.* (2003) 18:343–351. doi: 10.1359/jbmr.2003.18.2.343
67. Dukas L, Bischoff HA, Lindpaintner LS, Schacht E, Birkner-Binder D, Damm TN, et al. Alfacalcidol reduces the number of fallers in a community-dwelling elderly population with a minimum calcium intake of more than 500 mg daily. *J Am Geriatr Soc.* (2004) 52:230–6. doi: 10.1111/j.1532-5415.2004.52060.x
68. Chapuy MC, Arlot ME, Duboeuf F, Brun J, Crouzet B, Arnard S, et al. Vitamin D3 and calcium to prevent hip fractures in elderly women. *N Engl J Med.* (1992) 327:1637–42. doi: 10.1056/NEJM199212033272305
69. Smith H, Anderson F, Raphael H, Maslin P, Crozier S, Cooper C. Effect of annual intramuscular vitamin D on fracture risk in elderly men and women - a population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology.* (2007) 46:1852–7. doi: 10.1093/rheumatology/kem240
70. Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *J Am Med Assoc.* (2010) 303:1815–22. doi: 10.1001/jama.2010.594
71. Bischoff-Ferrari HA, Dawson-Hughes B, John Orav E, Staehelin HB, Meyer OW, Theiler R, et al. Monthly high-dose Vitamin D treatment for the prevention of functional decline a randomized clinical trial. *JAMA Intern Med.* (2016) 176:175–83. doi: 10.1001/jamainternmed.2015.7148
72. Smith LM, Gallagher JC, Suiter C. Medium doses of daily vitamin D decrease falls and higher doses of daily vitamin D3 increase falls: a randomized clinical trial. *J Steroid Biochem Mol Biol.* (2017) 173:317–22. doi: 10.1016/j.jsbmb.2017.03.015
73. Zhang HJ, Zhang JR, Mao CJ, Li K, Wang F, Chen J, et al. Relationship between 25-Hydroxyvitamin D, bone density, and Parkinson's disease symptoms. *Acta Neurol Scand.* (2019) 140:274–80. doi: 10.1111/ane.13141
74. Peterson AL, Mancini M, Horak FB. The relationship between balance control and vitamin D in Parkinson's disease—a pilot study. *Mov Disord.* (2013) 28:1133–7. doi: 10.1002/mds.25405

75. Hiller AL, Murchison CF, Lobb BM, O'Connor S, O'Connor M, Quinn JF. A randomized, controlled pilot study of the effects of vitamin D supplementation on balance in Parkinson's disease: does age matter? *PLoS ONE*. (2018) 13:e3637. doi: 10.1371/journal.pone.0203637
76. John EM, Schwartz GG, Koo J, Van Den Berg D, Ingles SA. Sun exposure, vitamin D receptor gene polymorphisms, and risk of advanced prostate cancer. *Cancer Res*. (2005) 65:5470–9. doi: 10.1158/0008-5472.CAN-04-3134
77. Gatto NM, Paul KC, Sinsheimer JS, Bronstein JM, Bordelon Y, Rausch R, et al. Vitamin D receptor gene polymorphisms and cognitive decline in Parkinson's disease. *J Neurol Sci*. (2016) 370:100–6. doi: 10.1016/j.jns.2016.09.013
78. Wilkins CH, Sheline YI, Roe CM, Birge SJ, Morris JC. Vitamin D deficiency is associated with low mood and worse cognitive performance in older adults. *Am J Geriatr Psychiatry*. (2006) 14:1032–40. doi: 10.1097/01.JGP.0000240986.74642.7c
79. May HT, Bair TL, Lappé DL, Anderson JL, Horne BD, Carlquist JF, et al. Association of vitamin D levels with incident depression among a general cardiovascular population. *Am Heart J*. (2010) 159:1037–43. doi: 10.1016/j.ahj.2010.03.017
80. Chan R, Chan D, Woo J, Ohlsson C, Mellström D, Kwok T, et al. Association between serum 25-hydroxyvitamin D and psychological health in older Chinese men in a cohort study. *J Affect Disord*. (2011) 130:251–9. doi: 10.1016/j.jad.2010.10.029
81. Pan A, Lu L, Franco OH, Yu Z, Li H, Lin X. Association between depressive symptoms and 25-hydroxyvitamin D in middle-aged and elderly Chinese. *J Affect Disord*. (2009) 118:240–3. doi: 10.1016/j.jad.2009.02.002
82. Anglin RES, Samaan Z, Walter SD, Sarah DM. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *Br J Psychiatry*. (2013) 202:100–7. doi: 10.1192/bjp.bp.111.106666
83. Jorde R, Sneve M, Figenschau Y, Svartberg J, Waterloo K. Effects of vitamin D supplementation on symptoms of depression in overweight and obese subjects: randomized double blind trial. *J Intern Med*. (2008) 264:599–609. doi: 10.1111/j.1365-2796.2008.02008.x
84. Penckofer S, Byrn M, Adams W, Emanuele MA, Mumby P, Kouba J, et al. Vitamin D Supplementation improves mood in women with type 2 diabetes. *J Diabetes Res*. (2017) 2017:8232863. doi: 10.1155/2017/8232863
85. Duval GT, Brangier A, Barré J, Launay CP, Beauchet O, Annweiler C. Vitamin D deficiency and incident onset of orthostatic hypotension in older adults: preliminary results from the “MERE” study. *J Am Geriatr Soc*. (2015) 63:1245–7. doi: 10.1111/jgs.13504
86. McCarroll KG, Robinson DJ, Coughlan A, Healy M, Kenny RA, Cunningham C. Vitamin D and orthostatic hypotension. *Age Ageing*. (2012) 41:810–3. doi: 10.1093/ageing/afs088
87. Ometto E, Stubbs B, Annweiler C, Duval GT, Jang W, Kim HT, et al. Hypovitaminosis D and orthostatic hypotension: a systematic review and meta-analysis. *J Hypertens*. (2016) 34:1036–43. doi: 10.1097/HJH.0000000000000907
88. Annweiler C, Schott AM, Rolland Y, Beauchet O. Vitamin D deficiency is associated with orthostatic hypotension in oldest-old women. *J Intern Med*. (2014) 276:285–95. doi: 10.1111/joim.12201
89. Soysal P, Yay A, Isik AT. Does vitamin D deficiency increase orthostatic hypotension risk in the elderly patients? *Arch Gerontol Geriatr*. (2014) 59:74–7. doi: 10.1016/j.archger.2014.03.008
90. Witham MD, Price RJG, Struthers AD, Donnan PT, Messow M, McConnachie A, et al. Effect of vitamin D supplementation on orthostatic hypotension: data from the Vitamin D in Isolated systolic hypertension randomized controlled trial. *J Hypertens*. (2014) 32:1693–9. doi: 10.1097/HJH.0000000000000223
91. Kim JE, Oh E, Park J, Youn J, Kim JS, Jang W. Serum 25-hydroxyvitamin D3 level may be associated with olfactory dysfunction in *de novo* Parkinson's disease. *J Clin Neurosci*. (2018) 57:131–5. doi: 10.1016/j.jocn.2018.08.003
92. Ross GW, Petrovitch H, Abbott RD. Serum vitamin D and risk of Parkinson's disease. *Mov Disord*. (2016) 31:935–93. doi: 10.1002/mds.26639
93. Ascherio A, Schwarzschild MA. The epidemiology of Parkinson's disease: risk factors and prevention. *Lancet Neurol*. (2016) 15:1257–72. doi: 10.1016/S1474-4422(16)30230-7
94. Suzuki M, Yoshioka M, Hashimoto M, Murakami M, Noya M, Takahashi D, et al. Randomized, double-blind, placebo-controlled trial of vitamin D supplementation in Parkinson disease. *Am J Clin Nutr*. (2013) 97:1004–13. doi: 10.3945/ajcn.112.051664
95. Habibi AH, Anamoradi A, Shahidi GA, Razmeh S, Alizadeh E, Kokhedan KM. Treatment of levodopa-induced dyskinesia with vitamin D: a randomized, double-blind, placebo-controlled trial. *Neurol Int*. (2018) 10:7737. doi: 10.4081/ni.2018.7737
96. Hollis BW. Circulating 25-Hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr*. (2005) 135:317–22. doi: 10.1093/jn/135.2.317
97. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. (2011) 96:1911–30. doi: 10.1210/jc.2011-0385
98. Lee JP, Tansey M, Jetton JG, Krasowski MD. Vitamin D toxicity: a 16-year retrospective study at an academic medical center. *Lab Med*. (2018) 49:123–9. doi: 10.1093/labmed/lmx077
99. Forsaa EB, Larsen JP, Wentzel-Larsen T, Herlofson K, Alves G. Predictors and course of health-related quality of life in Parkinson's disease. *Mov Disord*. (2008) 23:1420–7. doi: 10.1002/mds.22121
100. Thurman DJ, Stevens JA, Rao JK. Practice parameter: assessing patients in a neurology practice for risk of falls (an evidence-based review): report of the quality standards Subcommittee of the American Academy of Neurology. *Neurology*. (2008) 70:473–9. doi: 10.1212/01.wnl.0000299085.18976.20
101. Wood BH, Bilclough JA, Bowron A, Walker RW. Incidence and prediction of falls in Parkinson's disease: a prospective multidisciplinary study. *J Neurol Neurosurg Psychiatry*. (2002) 72:721–5. doi: 10.1136/jnnp.72.6.721
102. Bloem BR, Hausdorff JM, Visser JE, Giladi N. Falls and freezing of gait in Parkinson's disease: a review of two interconnected, episodic phenomena. *Mov Disord*. (2004) 19:871–84. doi: 10.1002/mds.20115
103. Bloem BR, van Vugt JP, Beckley DJ. Postural instability and falls in Parkinson's disease. *Adv Neurol*. (2001) 87:209–23.
104. Crouse JJ, Phillips JR, Jahanshahi M, Moustafa AA. Postural instability and falls in Parkinson's disease. *Rev Neurosci*. (2016) 27:549–55. doi: 10.1515/revneuro-2016-0002
105. Allen NE, Schwarzel AK, Canning CG. Recurrent falls in Parkinson's disease: a systematic review. *Parkinsons Dis*. (2013) 2013:906274. doi: 10.1155/2013/906274
106. Alexander NB. Gait disorders in older adults. *J Am Geriatr Soc*. (1996) 44:434–51. doi: 10.1111/j.1532-5415.1996.tb06417.x
107. Tinetti ME. Preventing falls in elderly persons. *N Engl J Med*. (2003) 348:42–9. doi: 10.1056/NEJMc020719
108. Dargent-Molina P, Favier F, Grandjean H, Baudoin C, Schott AM, Hausherr E, et al. Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet*. (1996) 348:145–9. doi: 10.1016/S0140-6736(96)01440-7
109. van der Marck MA, Klok MPC, Okun MS, Giladi N, Munneke M, Bloem BR, et al. Consensus-based clinical practice recommendations for the examination and management of falls in patients with Parkinson's disease. *Park Relat Disord*. (2014) 20:360–9. doi: 10.1016/j.parkrel.2013.10.030
110. Anastasiou CA, Yannakoulia M, Scarmeas N. Vitamin D and cognition: an update of the current evidence. *J Alzheimers Dis*. (2014) 42:S71–80. doi: 10.3233/JAD-132636
111. Llewellyn DJ, Langa KM, Lang IA. Serum 25-hydroxyvitamin D concentration and cognitive impairment. *J Geriatr Psychiatry Neurol*. (2009) 22:188–95. doi: 10.1177/0891988708327888
112. Llewellyn DJ, Lang IA, Langa KM, Muniz-Terrera G, Phillips CL, Cherubini A, et al. Vitamin D and risk of cognitive decline in elderly persons. *Arch Intern Med*. (2010) 170:1135–41. doi: 10.1001/archinternmed.2010.173
113. Peterson A, Mattek N, Clemons A, Bowman GL, Buracchio T, Kaye J, et al. Serum vitamin d concentrations are associated with falling and cognitive function in older adults. *J Nutr Heal Aging*. (2012) 16:898–901. doi: 10.1007/s12603-012-0378-4
114. Annweiler C, Schott AM, Allali G, Bridenbaugh SA, Kressig RW, Allain P, et al. Association of vitamin D deficiency with cognitive impairment in older women: cross-sectional study. *Neurology*. (2010) 74:27–32. doi: 10.1212/WNL.0b013e3181beecd3

115. Annweiler C, Rolland Y, Schott AM, Blain H, Vellas B, Herrmann FR, et al. Higher vitamin D dietary intake is associated with lower risk of Alzheimer's disease: a 7-year follow-up. *J Gerontol Ser A Biol Sci Med Sci.* (2012) 67:1205–11. doi: 10.1093/gerona/gls107
116. Corless D, Dawson E, Fraser F, Ellis M, Evans SJW, Perry JD, et al. Do vitamin D supplements improve the physical capabilities of elderly hospital patients? *Age Ageing.* (1985) 14:76–84. doi: 10.1093/ageing/14.2.76
117. Corless D, Ellis M, Dawson E, Fraser F, Evans S, Perry JD, et al. Using activities of daily living assessments to measure the effectiveness of vitamin d supplements in elderly long-stay patients. *Br J Occup Ther.* (1987) 50:60–2. doi: 10.1177/030802268705000208
118. Stein MS, Scherer SC, Ladd KS, Harrison LC. A randomized controlled trial of high-dose vitamin D2 followed by intranasal insulin in Alzheimer's disease. *J Alzheimer's Dis.* (2011) 26:477–484. doi: 10.3233/JAD-2011-110149
119. Rossom RC, Espeland MA, Manson JE, Dysken MW, Johnson KC, Lane DS, et al. Calcium and vitamin D supplementation and cognitive impairment in the women's health initiative. *J Am Geriatr Soc.* (2012) 60:2197–205. doi: 10.1111/jgs.12032
120. Masoumi A, Goldenson B, Ghirmai S, Avagyan H, Zaghi J, Abel K, et al. $1\alpha,25$ -dihydroxyvitamin D3 interacts with curcuminoids to stimulate amyloid- β clearance by macrophages of Alzheimer's disease patients. *J Alzheimers Dis.* (2009) 17:703–17. doi: 10.3233/JAD-2009-1080
121. Mizwicki MT, Liu G, Fiala M, Magpantay L, Sayre J, Siani A, et al. $1\alpha,25$ -dihydroxyvitamin D 3 and resolvin D1 return the balance between amyloid- β phagocytosis and inflammation in Alzheimer's disease patients. *J Alzheimers Dis.* (2013) 34:155–70. doi: 10.3233/JAD-121735
122. Langub MC, Herman JP, Malluche HH, Koszewski NJ. Evidence of functional vitamin D receptors in rat hippocampus. *Neuroscience.* (2001) 104:49–56. doi: 10.1016/S0306-4522(01)00049-5
123. Gezen-Ak D, Dursun E, Yilmazer S. Vitamin D inquiry in hippocampal neurons: consequences of vitamin D-VDR pathway disruption on calcium channel and the vitamin D requirement. *Neurol Sci.* (2013) 34:1453–8. doi: 10.1007/s10072-012-1268-6
124. Liang Q, Cai C, Duan D, Hu X, Hua W, Jiang P, et al. Postnatal vitamin D intake modulates hippocampal learning and memory in adult mice. *Front Neurosci.* (2018) 12:141. doi: 10.3389/fnins.2018.00141
125. Wang TT, Tavera-Mendoza LE, Laperriere D, Libby E, MacLeod NB, Nagai Y, et al. Large-scale *in silico* and microarray-based identification of direct $1,25$ -dihydroxyvitamin D3 target genes. *Mol Endocrinol.* (2005) 19:2685–95. doi: 10.1210/me.2005-0106
126. Schatz IJ, Bannister R, Freeman RL, Jankovic J, Koller WC, Low PA, et al. Consensus statement on the definition of orthostatic hypotension, pure autonomic failure and multiple system atrophy. *Clin Auton Res.* (1996) 144:218–9.
127. McDonald C, Newton JL, Burn DJ. Orthostatic hypotension and cognitive impairment in Parkinson's disease: Causation or association? *Mov Disord.* (2016) 31:937–46. doi: 10.1002/mds.26632
128. Senard JM, Raï S, Lapeyre-Mestre M, Brefel C, Rascol O, Rascol A, et al. Prevalence of orthostatic hypotension in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* (1997) 63:584–9. doi: 10.1136/jnnp.63.5.584
129. Allcock LM, Ulyart K, Kenny RA, Burn DJ. Frequency of orthostatic hypotension in a community based cohort of patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry.* (2004) 75:1470–1. doi: 10.1136/jnnp.2003.029413
130. Jang W, Park J, Kim JS, Youn J, Oh E, Kwon KY, et al. Vitamin D deficiency in Parkinson's disease patients with orthostatic hypotension. *Acta Neurol Scand.* (2015) 132:242–50. doi: 10.1111/ane.12390
131. Laird EJ, McNicholas T, O'Halloran AM, Healy M, Molloy AM, Carey D, et al. Vitamin D status is not associated with orthostatic hypotension in older adults. *Hypertension.* (2019) 74:639–44. doi: 10.1161/HYPERTENSIONAHA.119.13064
132. Arici Duz O, Helvacı Yılmaz N. Nocturnal blood pressure changes in Parkinson's disease: correlation with autonomic dysfunction and vitamin D levels. *Acta Neurol Belg.* (2019) doi: 10.1007/s13760-019-01113-7
133. Veronese N, Bolzetta F, De Rui M, Zambon S, Corti MC, Musacchio E, et al. Serum 25-hydroxyvitamin D and orthostatic hypotension in old people: the pro.V.A. study. *Hypertension.* (2014) 64:481–6. doi: 10.1161/HYPERTENSIONAHA.114.03143
134. Fullard ME, Tran B, Xie SX, Toledo JB, Scordia C, Linder C, et al. Olfactory impairment predicts cognitive decline in early Parkinson's disease. *Park Relat Disord.* (2016) 25:45–51. doi: 10.1016/j.parkreldis.2016.02.013
135. Boesveldt S, Verbaan D, Knol DL, Visser M, van Rooden SM, van Hilten JJ, et al. A comparative study of odor identification and odor discrimination deficits in Parkinson's disease. *Mov Disord.* (2008) 23:1984–90. doi: 10.1002/mds.22155
136. Doty RL, Deems DA, Stellar S. Olfactory dysfunction in Parkinsonism: a general deficit unrelated to neurologic signs, disease stage, or disease duration. *Neurology.* (1988) 38:1237–44. doi: 10.1212/WNL.38.8.1237
137. Ponsen MM, Stoffers D, Booij J, Van Eck-Smit BLE, Wolters EC, Berendse HW. Idiopathic hyposmia as a preclinical sign of Parkinson's disease. *Ann Neurol.* (2004) 56:173–81. doi: 10.1002/ana.20160
138. Braak H, Del Tredici K, Rüb U, De Vos RAI, Jansen Steur ENH, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging.* (2003) 24:197–211. doi: 10.1016/S0197-4580(02)00065-9
139. Deluca GC, Kimball SM, Kolasinski J, Ramagopalan S V., Ebers GC. Review: the role of vitamin D in nervous system health and disease. *Neuropathol Appl Neurobiol.* (2013) 39:458–84. doi: 10.1111/nan.12020
140. Kesby JP, Eyles DW, Burne THJ, McGrath JJ. The effects of vitamin D on brain development and adult brain function. *Mol Cell Endocrinol.* (2011) 347:121–7. doi: 10.1016/j.mce.2011.05.014
141. Glaser SD, Veenstra TD, Jirikowski GF, Prüfer K. Distribution of $1,25$ -dihydroxyvitamin D3 receptor immunoreactivity in the rat olfactory system. *Cell Mol Neurobiol.* (1999) 19:613–24.
142. Bohnen NI, Albin RL. The cholinergic system and Parkinson disease. *Behav Brain Res.* (2011) 221:564–73. doi: 10.1016/j.bbr.2009.12.048
143. Tillerson JL, Caudle WM, Parent JM, Gong C, Schallert T, Miller GW. Olfactory discrimination deficits in mice lacking the dopamine transporter or the D2 dopamine receptor. *Behav Brain Res.* (2006) 172:97–105. doi: 10.1016/j.bbr.2006.04.025
144. Wrzosek M, Lukaszkiwicz J, Wrzosek M, Jakubczyk A, Matsumoto H, Piatkiewicz P, et al. Vitamin D and the central nervous system. *Pharmacol Rep.* (2013) 65:271–8. doi: 10.1016/S1734-1140(13)71003-X
145. Klimmeck D, Mayer U, Ungerer N, Warnken U, Schnölzer M, Frings S, et al. Calcium-signaling networks in olfactory receptor neurons. *Neuroscience.* (2008) 151:901–12. doi: 10.1016/j.neuroscience.2007.11.023

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Fullard and Duda. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.